

Online ISSN: 2682-2628
Print ISSN: 2682-261X

IJC CBR

INTERNATIONAL JOURNAL OF CANCER AND BIOMEDICAL RESEARCH

<https://jcbr.journals.ekb.eg>

Editor-in-chief

Prof. Mohamed Labib Salem, PhD

Relationship Between Locus of Control Belief and IL-6 in Breast Cancer Patients

Azza M. El-Amir, Mohamed L. Salem, Noha L. Sabry, Eman M.
EL-Baiomy, Nehal EL- Mashad



PUBLISHED BY

EACR EGYPTIAN ASSOCIATION
FOR CANCER RESEARCH

Since 2014

Relationship Between Locus of Control Belief and IL-6 in Breast Cancer Patients

Azza M. El-Amir¹, Mohamed L. Salem², Noha L. Sabry³, Eman M. EL-Baiomy¹, Nehal EL- Mashad⁴

¹Immunology Division, Faculty of Science, Cairo University, Egypt

²Immunology and Biotechnology Unit, Faculty of Science, Tanta University, Egypt

³Psychiatry Department, Faculty of Medicine, Cairo University, Egypt

⁴Oncology Department, Faculty of Medicine, Tanta University, Egypt

ABSTRACT

Recent studies indicated to the immune disturbances in cancer patients in comparison with healthy individuals. Several studies suggested that individuals with external locus of control (E-LOC) have more symptoms of depression. Other studies indicated to the effective role of psychological status on the immune system. The association between E-LOC and immunity in cancer patients it is not clear. Study the relation between Rotter internal-external locus of control (I-E LOC) and God health locus of control (GHLC) and interleukin-6 (IL-6) in breast cancer patients. The study included hundred breast cancer patients and fifty healthy controls. Patients and participators responded to I-E LOC and GHLC questionnaires. Peripheral blood samples were collected to measure gene expression of IL-6 by using real time PCR. Breast cancer patients showed higher GHLC which was associated with higher level of IL-6 as compared to control subjects. By contrast, a significant inverse correlation between GHLC and IL-6 was found. However, there was no relationship between I-E LOC and IL-6. The results suggest no significant effect of LOC belief according to life on immune response against cancer, but high GHLC belief correlate with decreased IL-6. Further studies on the relations between other types of LOC and immune response in breast cancer is recommended.

Keywords: locus of control, breast cancer, IL-6

Editor-in-Chief: Prof. M.L. Salem, PhD - Article DOI: 10.21608/jcbr.2017.34733

ARTICLE INFO

Article history

Received: September 02, 2017

Revised: October 10, 2017

Accepted: October 23, 2017

Correspondence to

Azza El-Amir

Faculty of Science,

Cairo University,

Cairo, Egypt

Email: azzaelamir@yahoo.com

Copyright

©2017 Azza M. El-Amir, Mohamed L. Salem, Noha L. Sabry, Eman M. EL-Baiomy, Nehal EL- Mashad. This is an Open Access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any format provided that the original work is properly cited.

INTRODUCTION

Cancer is one of the major leading causes of death all over the world (Jemal et al., 2008). It has a negative effect on immune system as it causes loss of T cell specific, non-specific recognition and altered natural killer (NK) cells. It also associates with functional inhibition of macrophages and dendritic cells as well as increases in the recruitment of T regulatory (Treg) cells, myeloid-derived suppressor cells (MDSC) and monocytes (Jewett and Tseng, 2011).

Cancer is not affected only by biological factors but also by psychological factors which are known to have significant impact on the immune responses to cancer (Saxton et al., 2014). With this regard, one study on breast cancer suggested that women with depression

symptoms showed suppressed immunity (Turner-cobb et al., 2000). High inflammatory cytokines are characteristic of both depression and cachexia. The effects of cachexia can be more deleterious than the tumor growth itself, where thirty % of all patients with cancer die as a result of cachexia (Pirisi, 2000).

Previous studies have suggested a relationship between locus of control and depression. For instance, one -study indicated to an overall inverse relationship between spiritual well-being and depression has been reported (Holt et al., 2003). Another study also suggested that, awareness of God is associated with a better health (Ryan and Francis, 2012). Specifically, external locus of control (E. LOC) belief is often related to more symptoms of depression than internal locus of control (I-LOC) (Gibson et al., 2013). Accordingly, it can be suggested that

there is a relation between LOC belief and immune status in cancer patients. This study was aimed to investigate the relationship between I-E LOC and GHLC belief with IL-6 in breast cancer patients as compared to control subjects.

MATERIALS AND METHODS

Participants included 100 females (40-60 years old) with breast tumor at different stages. Participants were recruited from Damanhur Cancer Center and Tanta University Hospital, Egypt. Exclusion criteria included neurological disorders, ill conditions that affect immune status, diagnoses with another type of cancer, any anti-cancer treatment (chemotherapy, radiation, surgery) or any drugs that affect the immune system. Control subjects included 50 healthy females (40 to 60 years old), who were not diagnosed for any illness condition that affects the immune status or any psychological disorders. The research was approved by the ethical committee at Cairo University, Egypt.

Rotter internal-external locus of control (I-E LOC) scale

Rotter I-E LOC scale was used to measure LOC belief according to general life. It differentiates participants on one axis (internality-externality). High degree in I-E LOC means high E-LOC and low degree means high I-LOC (Rotter, 1966).

Questionnaire of God health locus of control (GHLC)

GHLC consists of 6 questions, which measures to what extent individual believes that God controls health. It is considered as a type of E-LOC. Each question ranged from 1= strongly disagree to 6= strongly agree. The total score ranges from 6 to 36 (Wallston et al., 1999).

Gene expression assay

RNA extraction was done by the Gene JETTM whole blood RNA purification mini kit. RNA was quantified in a single step assay by thermo scientific verso SYBER Green 1-step QRT-PCR kit plus ROX vial. Briefly, 500 µl of blood samples was centrifuged, the supernatant was discarded, and the pellet was lysed in 600µl lysis buffer. The lysate cells were washed by washing buffers. Then RNA was dissolved in RNA free

water and stored at -80°C until use. Master mix was prepared by adding enzyme mix, 1-step QPCR SYBER mix, RT enhancer, forward primer and reverse primer of IL-6 (Table 1). The cycling conditions were cDNA synthesis at 50°C for 15 min, thermo-start activation at 95°C for 15 min, denaturation at 95°C for 15 min, annealing at 95°C for 15 min and extension at 72°C for 30 sec. The relative level of mRNA expression of IL-6 was calculated based on $\Delta\Delta CT$ method and normalized to mRNA level of the house keeping gene $\beta 2$ macroglobulin.

Statistical analysis

Statistical presentation and analysis of the present study was analyzed by one-way ANOVA and bivariate co-relation by SPSS V.20.

Correlation between locus of control and pro-inflammatory cytokines in breast cancer

We then analyzed the correlation between locus of control belief and IL-6 in breast cancer patients. The results showed that significant inverse correlation between GHLC and IL-6 (Table 4) of control, I-E LOC: internal-external locus of control.

RESULTS

Locus of control belief in healthy and cancer breast patients.

This research studied Rotter I-E LOC and GHLC in healthy control and cancer breast patients. The results showed that patients had significantly higher GHLC than control subjects (Table 2), while they did not significantly different with regard to I-E LOC.

IL-6 in healthy control and cancer breast patients

We then analyzed IL-6 expression in patients with breast cancer and control subjects. The results showed that patients had significant higher level of IL-6 than controls (Table 3).

DISCUSSION

In the present study the LOC of breast cancer patients was correlated to the level of IL-6. LOC belief was assessed by I-E LOC and GHLC scales. The gene expression of IL-6 was assessed by real time PCR. Both I-E LOC and GHLC belief and IL-6 were studied in breast cancer patients and compared to control subjects.

Table 1. Primer used in measuring gene expression

| Gene | Sequence |
|----------------------------|--------------|
| B2 macroglobulin (Forward) | CCCCCACTGAA |
| B2-Microglobulin (Reverse) | TCATCCAATCCA |
| IL-6 (Forward) | GCCAGAGCTGT |
| IL-6 (Reverse) | AGGAACTCCT |

Table 2. Locus of control belief in healthy control and cancer breast patient's groups

| Group | GHLC | I-E LOC |
|----------|---------------|-------------|
| Controls | 26.69 ± 0.92 | 8.44 ± 0.58 |
| Patients | 32.13 ± 1.15* | 9.63 ± 0.38 |

* Significantly different as compared to another group ($p < 0.05$).

Table 3. Pro-inflammatory cytokines in healthy control and cancer breast patients

| Group | IL-6 |
|----------|--------------|
| Controls | 1.00 |
| Patients | 2.19 ± 0.26* |

* Significantly different as compared to control ($p < 0.05$).

Table 4. Correlation between locus of control and IL-6 in breast cancer patients' group

| Locus type | Correlation coefficient | IL-6 |
|------------|-------------------------|--------|
| I-E LOC | r | -0.38 |
| GHLC | r | -0.67* |

* Significant at $p \leq 0.05$. GHLC: God health locus

When healthy controls and breast cancer patients were compared, no significant difference was found with regard to I-E LOC between patients and controls. However, it was observed that patients showed higher GHLC than healthy controls. Similar finding was concluded in a previous study which suggested that women with breast cancer are higher GHLC belief than healthy women (Iskandarsyah et al., 2014). These data suggest that facing serious diseases such as cancer call for increasing needs for external support.

Regarding to IL-6, it was found that patients express higher serum levels of IL-6 than healthy controls. These results are consistent with a previous study, which suggested that IL-6 is dramatically up-regulated in breast cancer patients in comparison with healthy controls (Nariřa et al., 2011).

In addition, another study on breast cancer patients indicated that high levels of IL-6 is associated with poor overall survival (ShuChen et al., 2015).

By correlating the relationship between IL-6 and LOC, it was found that there is no significant correlation between I-E LOC and IL-6. By contrast, there was a significant inverse correlation between GHLC and IL-6. These data would suggest the external support to cancer patients may require adjuvant therapy to correct the balance in immune system. It also suggests that GHLC belief itself is important for a healthy immune response in breast cancer since IL-6 was found to be associated with poor clinical outcomes in breast cancer patients (Ravishankaran and Karunanithi, 2011). To the best of our knowledge, the present study is the first to investigate the relationship between GHLC belief and IL-6.

In conclusion, although there was no significant difference in LOC belief according to general life between healthy control and patients, cancer patients expressed higher belief in God belief as a controller of their health than healthy controls. Also, patients had higher IL-6 expression than healthy controls. As such, both IL-6 and GHLC belief might be important factors in response of cancer patients to combat this disease.

ACKNOWLEDGMENT

This study was partially supported by a grant from Science and Technology Development Fund (STDF), Ministry of Higher Education and Scientific Research, Egypt to Mohamed L Salem. We are grateful to all breast cancer patients and volunteers who took part in the present research study.

REFERENCES

- Gibson RC, Morgan KA, Abel WD, Sewell CA, Martin JS, Lowe GA, Haye Wde L, Edwards CL, O'Garro KN, Reid ME and Asnani MR (2013). Locus of control, depression and quality of life among persons with sickle cell disease in Jamaica. *Psychology, Health and Medicine*; 18: 451-460.
- Holt CL, Clark EM, Kreuter MW and Rubio DM (2003). Spiritual health locus of control and breast cancer beliefs among urban African American women. *Health Psychology*; 22: 294-299.
- Iskandarsyah A, de Klerk C, Suardi DR, Sadarjoen SS and Passchier J (2014). Health locus of control in Indonesian women with breast cancer: a comparison with healthy women. *Asian Pacific Journal of Cancer Prevention*; 15: 9191-9197.

- Jemal A, Siegel R, Ward E, Hao Y, XU J, Murray T and Thun Mj (2008). Cancer statistics. CA: a Cancer Journal for Clinicians; 58: 71-96.
- Jewett A and Tseng HC (2011). Tumor induced inactivation of natural killer cell cytotoxic function; implication in growth, expansion and differentiation of cancer stem cells. Journal of Cancer; 2: 443-457.
- Narița D, Seclaman E, Ursoniu S, Iliina R, Cireap N and Anghel A. (2011). Expression of CCL18 and interleukin-6 in the plasma of breast cancer patients as compared with benign tumor patients and healthy controls. Romanian Journal of Morphology and Embryology; 52: 1261-1267.
- Pirisi A (2000). US researchers find key link in muscle-wasting syndrome. The Lancet; 356: 9237: 1249.
- Ravishankaran P and Karunanithi R. (2011). Clinical significance of preoperative serum interleukin-6 and C-reactive protein level in breast cancer patients. World Journal of Surgical Oncology; 6: 9:18
- Rotter JB (1966). Generalized expectancies for internal versus external control of reinforcement. Psychol Monogr; 80: 1-28.
- Ryan ME and Francis AJ (2012). Locus of control beliefs mediate the relationship between religious functioning and psychological health. Journal of Religion and Health; 51: 774-785.
- Saxton JM, Scott EJ, Daley AJ, Woodrooffe M, Mutrie N, Crank H, Powers HJ and Coleman RE (2014). Effects of an exercise and hypocaloric healthy eating intervention on indices of psychological health status, hypothalamic- pituitary-adrenal axis regulation and immune function after early-stage breast cancer: a randomised controlled trial. Breast Cancer Research and Treatment; 16: R39.
- ShuChen Lin, ZhiHua Gan, Kun Han, Yang Yao and DaLiu Min (2015) Interleukin-6 as a prognostic marker for breast cancer: a meta-analysis. Tumori; 101: 535 – 541.
- Turner-cobb JM, Sephton SE, Koopman C, Blake-mortimer J and Spiegel D (2000). Social support and salivary cortisol in women with metastatic breast cancer. Psychosomatic Medicine; 62: 337–345.
- Wallston KA, Malcarnevl, Flores L, Hansdottir I, Smith CA, Stein MJ, Weisman MH and Clements PJ (1999). Does God determine your health? The God Locus of Control scale. Cognitive Therapy and Research.